Communications of BAO, Vol. 73, Issue 2, 2026, pp. 363-366

Radio properties of Blazars

V.Kh. Mkrtchyan *, H.V. Abrahamyan †, and A.M. Mickaelian‡

Byurakan Astrophysical Observatory

Abstract

Blazars represent the most extreme class of Active Galactic Nuclei (AGN), characterized by powerful relativistic jets aligned close to the line of sight of the observer. In this work, we present a comprehensive multi-frequency study of the radio properties of blazars extracted from the 5th edition of the Roma-BZCAT catalog. By performing precise cross-matching across 14 independent radio survey catalogs spanning frequencies from 74 MHz to 8400 MHz (including NVSS, FIRST, TGSS, LOFAR, WENSS, and GB6), we construct broadband radio spectral distributions for the different blazar subclasses. From a preliminary representative sample of 20 objects per subclass (BZB, BZG, BZQ, and BZU), we calculated the mean radio spectral indices (α). Our results demonstrate a clear physical distinction among the subclasses: Flat Spectrum Radio Quasars (BZQ) exhibit a predominantly flat spectral index (α_{avg} = -0.43), driven by compact, self-absorbed core jet emission, whereas Blazar-like Galaxies (BZG) show significantly steeper radio spectra (α_{avg} = -0.93), indicating major synchrotron contributions from extended radio lobes. Furthermore, we outline our ongoing framework for combining FIRST radio fluxes with SDSS optical photometry to evaluate radio-to-optical flux ratios and their correlations across blazar populations.

Keywords: *Blazars, Active Galactic Nuclei (AGN), BZCAT, Radio Spectral Index, Relativistic Jets*

1. Introduction

Active Galactic Nuclei (AGN) are among the most luminous and energetic phenomena in the universe, powered by accretion onto supermassive black holes ($M \sim 10^6 - 10^9\, M_\odot$) located at the centers of distant host galaxies. Among the diverse zoo of AGN, blazars constitute an extreme subclass where a relativistic, non-thermal jet of plasma is oriented at a very small angle ($\theta \leq 10^\circ$) relative to the observer's line of sight.

Due to Doppler boosting and relativistic beaming, the observed radiation from blazars is strongly amplified, resulting in:

- High luminosity across the entire electromagnetic spectrum (from radio to high-energy γ-rays).
- Extreme and rapid temporal variability (resembling cosmic fireworks).
- High linear polarization in both optical and radio bands.

Blazars are traditionally classified into two primary physical categories based on their optical spectral properties: BL Lacertae (BL Lac) objects, which feature featureless optical continuum spectra or very weak emission lines, and Flat Spectrum Radio Quasars (FSRQs), which display broad, strong optical emission lines.

Understanding the radio spectral behavior of these sub-classes is paramount for constraining jet dynamics, synchrotron self-absorption mechanisms, energy dissipation regimes, and the interaction between the central engine and the extended environment. The primary goal of this study is to systematically cross-match the latest blazar catalogs with multi-frequency radio surveys, calculate their radio spectral indices (α), and establish the connection between radio emission profiles, optical properties, and blazar taxonomy.

*varduhi.mkrtchyan.99@bk.ru, Corresponding author
†abrahamyanhayk@gmail.com
‡aregmick@yahoo.com

doi: https://doi.org/10.52526/25792776-26.73.2-363

2. Sample and Catalog Selection

2.1. The Roma–BZCAT Catalog

The primary target list was derived from the 5th edition of the Multi-Frequency Catalog of Blazars (Roma-BZCAT, Massaro et al.), which contains a total of 3,561 objects. The catalog categorizes sources into four major subclasses based on optical spectral energy distributions (SED) and line characteristics:

- **5BZB (BL Lac Objects):** 1,151 sources (32.3% of the catalog) — AGNs with featureless optical spectra or showing only weak absorption lines of galactic origin.
- **5BZG (Galaxies with Blazar Activity):** 274 sources (7.7%) — Sources usually reported as BL Lacs in the literature, but whose SED displays a strong thermal contribution from the host galaxy over the nuclear emission.
- **5BZQ (Flat Spectrum Radio Quasars):** 1,909 sources (53.6% of the catalog) — Luminous quasars exhibiting broad optical emission lines and prominent blazar features.
- **5BZU (Unclassified Blazars):** 227 sources (6.4%) — Objects exhibiting transition features, peculiar broad-line variability, or uncertain blazar characteristics.

2.2. The Roma–BZCAT Catalog

To analyze the radio continuum properties over two orders of magnitude in frequency, we cross-matched the BZCAT coordinates against 14 major radio survey catalogs using tailored cross-matching radii optimized for each survey's spatial resolution:

Catalog Name	Survey Frequency	Angular Resolution	Search Radius	Matched Sources
VLSSr	74 MHz	∼80”	45”	937
LOFAR (LoTSS)	120–168 MHz (144 MHz)	∼6”	15”	956
TGSS ADR1	150 MHz	∼25”	45”	2,656
7C	151 MHz	∼70”	45”	514
Miyun	232 MHz	∼150”	60”	394
WENSS	330 MHz	∼54”	45”	1,024
Texas	365 MHz	∼50”	60”	1,182
Molonglo	408 MHz	∼163”	60”	10
SUMSS	843 MHz	∼45”	45”	545
FIRST	1400 MHz (1.4 GHz)	∼5”	15”	1,650
NVSS	1400 MHz (1.4 GHz)	∼45”	45”	3,186
GB6	4850 MHz	∼210”	45”	1,700
87GB	4850 MHz	∼210”	45”	1,541
CLASS	8400 MHz	∼0.2”	60”	1,583

2.3. Focus on Key 1.4 GHz Surveys: FIRST vs. NVSS

Two major VLA surveys at 1.4 GHz provide fundamental data for our morphological and flux density studies:

- **FIRST (Faint Images of the Radio Sky at Twenty-Centimeters):** Covers ∼10,000 deg^2 of the northern galactic cap with a high angular resolution of ∼5” and a sensitivity threshold of ∼1 mJy. It provides detailed morphological structures of blazar cores and jets.
- **NVSS (NRAO VLA Sky Survey):** Covers ∼82% of the sky north of $\delta = -40°$ (∼35,000 deg^2) with a synthesis resolution of ∼45”. NVSS is essential for capturing total integrated radio fluxes, including diffuse, large-scale structures.

3. Methodology & Spectral Index Calculations

The radio spectral index α is defined according to the power-law relation between flux density S_ν and observing frequency ν:

$$S_\nu \propto \nu^\alpha \implies \log(S_\nu) = \alpha \cdot \log(\nu) + C$$

In the preliminary stage of this project, a representative subset of 20 objects per blazar subclass was selected. For each object, multi-frequency flux measurements were compiled, and a linear regression fit in $\log(S_\nu) - \log(\nu)$ space was executed to determine the individual radio spectral indices.

4. Preliminary Results and Discussion

The calculated average radio spectral indices (α_{avg}) for each blazar subclass are summarized in the table below:

Subclass	Source Description	Average Spectral Index	Dominant Spectral Behavior
5BZQ	Flat Spectrum Radio Quasars	**−0.43**	Flat
5BZB	BL Lacertae Objects	**−0.78**	Steep
5BZU	Unclassified Blazars	**−0.83**	Steep
5BZG	Blazar-like Galaxies	**−0.93**	Steep

Physical Interpretation of Findings:

- **Flat Spectrum Radio Quasars (BZQ):** BZQs exhibit the flattest mean radio spectrum ($\alpha = -0.43$). This flat profile ($\alpha > -0.5$) is a classical signature of continuous synchrotron self-absorption occurring near the optical core of a highly beamed, compact relativistic jet. The superposition of multiple compact jet components, each peaking at a different frequency, creates a flat composite spectrum.
- **Blazar-like Galaxies (BZG):** BZGs show the steepest radio spectra ($\alpha = -0.93$). In these systems, the Doppler-boosted core emission is less dominant relative to the extended, unbeamed radio lobes. The extended lobes emit optically thin synchrotron radiation, which naturally produces steeper spectral slopes due to radiative cooling of aging electron populations.
- **BL Lac Objects (BZB) & Unclassified Sources (BZU):** BZB ($\alpha = -0.78$) and BZU ($\alpha = -0.83$) sources demonstrate intermediate spectral slopes. This suggests a hybrid emission state where both compact core jets and extended lobe structures contribute significantly to the total integrated flux density.



Figure 1. Radio spectrum examples of blazar types.

doi: https://doi.org/10.52526/25792776-26.73.2-363

5. Conclusions and Future Work

Our initial multi-frequency cross-matching and spectral analysis confirm distinct radio spectral behaviors among blazar subclasses, providing valuable insight into their underlying physics and beaming geometry.

Immediate and Future Objectives:

- **Full Sample Spectral Fitting:** Extend the spectral index (α) calculation to all 3,561 BZCAT sources by applying automated robust linear and parabolic spectral fitting algorithms.
- **Radio-to-Optical Ratio Determination:** Calculate the broad-band radio-to-optical flux ratio (R_{ratio}), utilizing 1.4 GHz FIRST radio fluxes and optical g, r, i, z magnitudes from the Sloan Digital Sky Survey (SDSS).
- **Correlation Analyses:**
  - Investigate correlations between R_{ratio} and optical spectral properties across subclasses.
  - Analyze the relation between the radio spectral index α and the optical color/luminosity parameters.
  - Probe redshift-dependent evolution of radio spectral steepening across the BZQ and BZB populations.

References

Abrahamyan H. V., 2020, Astronomische Nachrichten, 341, 703

Abrahamyan H. V., 2021, Communications of the Byurakan Astrophysical Observatory, 68, 130

Abrahamyan H. V., Mickaelian A. M., Mikayelyan G. A., Paronyan G. M., 2025, Communications of the Byurakan Astrophysical Observatory, 72, 396

Abrahamyan H. V., Mickaelian A. M., Mikayelyan G. A., Paronyan G. M., Dennefeld M., 2026, Astrophysics,

Becker R. H., White R. L., Helfand D. J., 1995, Astrophys. J. , 450, 559

Condon J. J., Cotton W. D., Greisen E. W., Yin Q. F., Perley R. A., Taylor G. B., Broderick J. J., 1998, Astron. J. , 115, 1693

Massaro E., Maselli A., Leto C., Marchegiani P., Perri M., Giommi P., Piranomonte S., 2015, Astrophys. Space. Sci. , 357, 75

Mkrtchyan V. K., Abrahamyan H. V., Mickaelian A. M., 2025a, Communications of the Byurakan Astrophysical Observatory, 72, 100

Mkrtchyan V. K., Mickaelian A. M., Abrahamyan H. V., 2025b, Communications of the Byurakan Astrophysical Observatory, 72, 423

Paronyan G. M., Mickaelian A. M., Abrahamyan H. V., Mikayelyan G. A., 2021, Astrophysics, 64, 277